

On the Multifunctional Role of the Biota in the Self-Purification of Aquatic Ecosystems

S. A. Ostroumov

Moscow State University, Vorob'evy gory, Moscow, 119992 Russia

Received November 28, 2003

Abstract—Principles of the theory of the ecological mechanism of water self-purification based on multiple functions of the biota in freshwater and marine ecosystems are formulated. In developing this theory, the results of the author's experiments with filtering hydrobionts have been used. These results indicate that the water self-purification mechanism is vulnerable to the impact of some pollutants and, in particular, surfactants. Conclusions drawn on the basis of the theory have practical significance for biodiversity conservation and for the sustainable use of the biological resources of aquatic ecosystems.

Key words: aquatic ecosystems, water quality, water self-purification, pollution, theory.

The study of water self-purification processes (Skurlatov, 1988; Ostroumov, 2000a) is essential for approaching fundamental ecological problems (Alimov, 2000; Ostroumov *et al.*, 2003) and for resolving applied issues related to the sustainable use of natural resources.

The purpose of this paper is to give a systematic account of the concepts concerning the multiple functions of the biota in the self-purification of water bodies and watercourses, without attempting to review the numerous publications in this field. A quantitative assessment of the various processes involved in the self-purification of water bodies and watercourses is also beyond the scope of this study.

An analysis of our publications over the period from 1997 to 2005 allowed me to make some generalizations concerning the main processes and factors involved in the self-purification system of aquatic ecosystems.

MAIN PHYSICAL, CHEMICAL, AND BIOTIC PROCESSES LEADING TO WATER PURIFICATION IN AQUATIC ECOSYSTEMS

These processes are listed in Table 1. Some of them were considered in a number of publications (Alimov, 1981; Izrael' and Tsyban', 1989; Lisytsin, 2001; Matishov and Matishov, 2001).

Physical and chemical processes of water self-purification are often regulated by biotic factors or strongly depend on them (Ostroumov, 2001a, 2002c). Thus, the degree of pollutant sorption on settling suspended particles depends on the concentration of phytoplankton cells, photochemical processes depend on water transparency, and transparency depends on the filtering capacity of hydrobionts. Free-radical processes of pollutant destruction depend on the binding of metal ions

to dissolved ligands, namely, organic molecules of biological origin (Skurlatov, 1988). Thus, biotic factors are at the center of the entire water self-purification system.

All processes involved in water purification are important, and none of them can be considered more significant than others. The important biotic processes leading to water purification that have been characterized in detail include organic matter oxidation (Sadchikov, 1997; Zavarzin and Kolotilova, 2001; Wetzel, 2001) and water filtration by hydrobionts (Alimov, 1981; Sushchenya, 1975; Shul'man and Finenko, 1990).

The activity of a community that oxidizes organic matter can be expressed in absolute and relative values: for instance, as the ratio of energy expended by hydrobionts for metabolism (total respiration R) to their total biomass (B). The $(R/B)_e$ ratio, referred to as the Schrödinger ratio, reflects the relationship between energy expenditures for the maintenance of life activity and, thereby, structure of the community and the amount of energy contained in its structure (Alimov, 2000).

Organic matter is oxidized by many hydrobionts, with a special role belonging to bacteria (Vinogradov and Sushkina, 1987; Zavarzin and Kolotilova, 2001; Wetzel, 2001). The total biomass of bacteria in the epipelagic zone (0–200 m) of the World Ocean is approximately 276×10^6 t C, averaging 8 g of fresh biomass (0.8 g C) per square meter of water surface (Vinogradov and Shushkina, 1987). Many specialists consider that bacteria account for 60–70% of the total heterotrophic destruction in the ocean. However, estimations of global heterotrophic destruction in the ocean are ambiguous. According to calculations based on average O_2 consumption by one bacterial cell, bacterial destruction in the World Ocean amounts to 100×10^9 t C per year; in this

Table 1. Some factors and processes involved in water self-purification (PI, pollutants; numbers beginning with 1, 2, and 3 indicate physical, chemical, and biological factors and processes, respectively)

No.	Factors and processes of water purification	Biotic factors that affect the processes indicated on the left
1.1	Dissolution and dilution	Mixing depends on macrophytes
1.2	Transfer on land	The same
1.3	Transfer to adjoining water bodies	»
1.4	Sorption of PI by suspended particles followed by their sedimentation	Seston formation
1.5	Sorption of PI by bottom sediments	Detritus formation
1.6	Evaporation	Properties of the surface film depend on DOM
2.1	Hydrolysis	pH depends on photosynthesis
2.2	Photochemical transformation	DOM and suspended organic matter
2.3	Catalytic redox transformations	DOM
2.4	Transformations with the involvement of free radicals	DOM
2.5	Decrease in PI toxicity resulting from binding to dissolved organic matter (DOM)	DOM
2.6	Chemical oxidation of PI with the involvement of oxygen	Photosynthesis
3.1	Release of oxygen that oxidizes PI	Photosynthesis
3.2	Sorption and accumulation of PI and biogenic substances by hydrobionts	Plankton, benthos
3.3	Biotransformation of PI	Enzymes
3.4	Extracellular enzymatic transformation of PI	Exoenzymes
3.5	Removal of suspended particles from water by filtration	Filter feeders
3.6	Removal of PI from water column as a result of sorption by pellets	Filter feeders
3.7	Release into water of organic substances that serve as sensitizers promoting PI photolysis	DOM
3.8	Release into water of organic substances binding with PI to form less toxic complexes	DOM
3.9	Release into water of substances involved in free-radical and catalytic redox mechanisms of PI destruction	DOM
3.10	Prevention or retardation of the release of biogenic substances and PI into water and their accumulation by benthic organisms	Benthos
3.11	Increase in organic matter content in bottom sediments, which improves PI binding by bottom sediments	Detritus formation
3.12	Release of N and P compounds into water (element recycling), which promotes the growth of oxygen-releasing phototrophic organisms	Bacteria of bottom sediments
3.13	Removal of C, N, and P from an aquatic ecosystem due to emergence of aquatic insects, migration of amphibians on land, and foraging of fish-eating birds	Biota of an ecosystem
3.14	PI biotransformation and sorption in soil (when polluted water is used for irrigation)	The same
3.15	Regulation of the abundance and activity of organisms involved in water purification	»

case, the entire heterotrophic destruction in the 0–200 m layer is 150×10^9 t C per year. In the variant based on the daily average specific production of bacteria and the coefficient characterizing the efficiency of assimilated food utilization for growth ($K_2 = 0.33$), bacterial destruction is approximately 60×10^9 t C per year, and total heterotrophic destruction is 85×10^9 t C per year. An important role in heterotrophic destruction belongs to protists (total biomass in the epipelagial 69×10^6 t C,

averaging 0.2 g C/m²) and metazooplankton (body size 0.2–5 mm; total biomass in the epipelagial 386×10^6 t C, averaging 1 g C/m²). The ratio of daily production to biomass (P/B) in the epipelagial averages 53% for protists and 2% for metazooplankton (Vinogradov and Sushkina, 1987).

The filtration rate in some groups of hydrobionts (ascidians, cirripeds, bryozoans, echinoderms, bivalves, gastropods, polychaetes, and sponges) often

ranges from 1 to 8.8 l/h per gram dry (decalcified) body weight (Dame *et al.*, 2001). The dependence of filtration rate on body weight is described by an exponential function (Alimov, 1981). The total water filtration by populations of macroinvertebrates (Mollusca, Ascidia, Polychaeta) in the water column above 1 m² of the bottom was estimated at 1–10 m³ per day (Ostroumov, 2001a). A great ecological significance of filtering hydrobionts was emphasized by K.A. Voskresenskii and A.S. Konstantinov (for review, see Ostroumov, 2001a).

MAIN FUNCTIONAL BLOCKS OF THE SELF-PURIFICATION SYSTEM IN AQUATIC ECOSYSTEMS

The main functional blocks accounting for the major part of the total hydrobiological mechanism of self-purification in aquatic ecosystems are as follows: (1) the block of filtering activity (“filters”) (Ostroumov, 1998); (2) the block of mechanisms accounting for the transfer (pumping) of chemical substances from one ecological compartment (environment) to another, i.e., “pumps” operating in the self-purification mechanisms of aquatic ecosystems; and (3) the block responsible for the cleavage of pollutant molecules (“mills” that crush pollutants).

Filters. Four filtering systems are distinguished (Ostroumov, 1998): (a) the aggregate of invertebrate filter-feeding hydrobionts (Ostroumov, 2001a; Dame *et al.*, 2001; Ostroumov, 2005); (b) the belt of coastal macrophytes that retains a portion of the biogenic substances and pollutants coming to the ecosystem from the adjoining territory; (c) the benthos that retains and absorbs some of the biogenic substances and pollutants migrating at the water–bottom sediment interface; and (d) microorganisms sorbed on suspended particles that move relative to the water mass (settle) because of gravity; as a result, microorganisms absorb dissolved organic substances and biogenic elements from the water (Ostroumov, 1998). During the sedimentation of a water-suspended particle, oxygen exchange between the bacteria sorbed on it and the aquatic environment increases (Zavarzin and Kolotilova, 2001).

Pumps. The following functional systems promote substance transfer from one site to another (Ostroumov, 2001a): (a) the block of processes acting as a pump that transfers some pollutants from the water column to sediments (e.g., sedimentation and sorption); (b) the block of processes acting as a functional pump that promotes the transfer of some pollutants from the water column to the atmosphere (evaporation); (c) the block of processes acting as a functional pump that promotes the transfer of some biogenic substances from water to the surrounding terrestrial ecosystems (the sum of migration processes related to the emergence of adult insects in the species whose larvae develop in water); (d) an analogous block of processes involving the transfer of

some biogenic substances from water to the surrounding terrestrial ecosystems due to the activities of birds that feed on hydrobionts, removing biomass from an aquatic ecosystem, but nest on the surrounding territory; and (e) the block of processes involving the transfer of some biogenic substances from water to the coastal ecosystems due to the emergence from a water body of amphibians whose early development proceeds in water.

Mills are the functional systems that destroy an excess of organic matter and cleave pollutants (Ostroumov, 1986, 2001a; Skurlatov, 1988): (a) the molecular mill of intracellular enzymatic processes, (b) the mill of extracellular enzymes present in an aquatic environment, (c) the mill of photochemical processes promoted by sensitizers of biological origin, and (d) the mill of free radical processes involving ligands of biological origin (Skurlatov, 1988). There are many examples of the quantitative study of the corresponding processes (Wetzel, 2001).

SOURCES OF ENERGY FOR THE BIOTIC MECHANISMS OF AQUATIC ECOSYSTEM SELF-PURIFICATION

The biotic processes of self-purification receive energy from the following sources: photosynthesis, oxidation of autochthonous organic matter, oxidation of allochthonous organic matter, and other redox reactions. Thus, virtually all available energy sources are used. Some energy is received due to oxidation of the components of which the system rids itself (dissolved and suspended organic matter). In this respect, self-purification processes are comparable to energy-saving technologies (Ostroumov, 2002b).

Self-purification is often related to organic matter oxidation by aerobic organisms. Equally important are anaerobic processes, in which energy is generated due to electron transfer to acceptors other than oxygen. The anaerobic energetics determines the metabolism of many microorganisms, including the methanogenic (destruction of organic matter with the release of methane), sulfidogenic (the release of H₂S, H₂, and methane), and anoxygenic phototrophic communities (the release of SO₄²⁻, H₂S, hydrogen, and methane) (Zavarzin and Kolotilova, 2001). The products of their activity are further used as oxidation substrates by organisms of other communities, including bacteria of the group named “the bacterial oxidative filter.” This group functions under aerobic conditions and oxidizes hydrogen (hydrogen-reducing bacteria), methane (methanophores), NH₃ (nitrifiers), H₂S (sulfur bacteria), and thiosulfate (thionic bacteria). The involvement of bacteria that use metal ions or ions containing metals (Fe, Mn) as electron acceptors is also important. Quantitative characteristics of relevant processes are given in the paper by Wetzel (2001).

INVOLVEMENT OF MAIN GROUPS OF ORGANISMS IN THE SELF-PURIFICATION OF AQUATIC ECOSYSTEMS

The involvement of microorganisms, phytoplankton, higher plants, invertebrates, and fish in the self-purification of aquatic ecosystems and the improvement of water quality was analyzed in several papers (Ostroumov, 1986, 2000c, 2001a; Zavarzin and Kolotilova, 2001). All these groups take an active part in the self-purification of aquatic ecosystems, each being involved in more than one or two processes.

Heterotrophic aerobic bacteria actively participate in self-purification processes. As noted above, representatives of many other groups of bacteria are also involved in organic matter destruction and self-purification of water bodies (Zavarzin and Kolotilova, 2001). Moreover, an important role in organic matter destruction belongs to protists, aquatic fungi, and related organisms.

The diversity of microorganisms participating in the destruction of biopolymers and in the self-purification system is the more important as the microorganisms that functionally complement each other in bacterial communities are represented by phylogenetically distant forms (Zavarzin and Kolotilova, 2001).

Water filtration, which is of primary importance for water purification, is accomplished by representatives of many taxa (Alimov, 1981; Sushchenya, 1975; Vinogradov and Shushkina, 1987; Ostroumov, 2005). In the marine plankton, for instance, the function of filter feeders consuming fine particles (nanophages) is performed by Appendiculariae, Doliolida (class Thaliacea), small Calanoida, meroplankton (larvae), and other invertebrates; the function of coarse filtration (euryphagous filter-feeders) is performed by *Oithona* sp. (Cyclopoida), Oncaea (Cyclopoida), large Calanoida, and Euphausiacea (Vinogradov and Sushkina, 1987). For a detailed list of the main planktonic and benthic filter feeders in ecosystems, see Ostroumov (2002a).

RELIABILITY OF THE WATER SELF-PURIFICATION SYSTEM

The reliability of the water self-purification mechanism is closely related to the fundamentally important problem of ecosystem stability (Krasnoshchekov and Rozenberg, 1992) and is often ensured by duplicating many components of the system. For instance, water filtration is performed by two different large groups of organisms, the plankton and the benthos, and at a high rate (Alimov, 1981; Sushchenya, 1975). In addition, the benthos duplicates the activity of zooplankters constantly remaining in the pelagial zone, because the larvae of many benthic filter feeders are planktonic organisms. The plankton includes two large groups of multicellular invertebrate filter feeders, crustaceans (Sushchenya, 1975), and rotifers (Monakov, 1998), which provide a backup for each other. In addition,

there is yet another large group of organisms (Protozoa) with a slightly different type of feeding, which, in turn, provide a backup for multicellular filter feeders (crustaceans and rotifers).

Water filtration is performed in parallel by many representatives of the biota. The filtering activity (the amount of water filtered per hour as expressed in body volumes of the filter-feeding organism) is up to 5×10^6 in nanoflagellates and 5×10^5 in ciliates (Fenchel, 1986, 1987; cited from Wetzel, 2001). Cladocerans filter 4–40 ml (in some cases, up to 130 ml) per animal per day; copepods, up to 27 ml (Wetzel, 2001); rotifers, 0.07–0.3 ml (Monakov, 1998). These and other filter feeders provide a backup for each other in removing suspensions from the water.

Another group of important self-purification processes, the enzymatic destruction of pollutants, are accomplished by bacteria and fungi, which, therefore, provide a backup for each other. The same applies to the variety of hydrobionts participating in the oxidation of dissolved organic matter.

Yet another important component of reliability is the self-regulation of the biota. Almost all organisms involved in self-purification are under the dual control of other organisms representing the preceding and subsequent links in the trophic chain. Their role can be efficiently studied by means of the inhibitory analysis of regulatory interactions in trophic chains (Ostroumov, 2000b).

Among the mechanisms of ecosystem regulation, an important role belongs to different forms of signaling, including those by means of chemical substances that transmit information or regulatory signals. I have proposed to designate such substances ecological chemoregulators and ecological chemomediators (Ostroumov, 1986, 2003).

The reliability of the water self-purification system in an ecosystem is provided by the multiplicity of components (processes) that comprise this system and operate, to some extent, in parallel. Water purification and the improvement of its quality, in turn, are necessary for the self-maintenance of the entire aquatic ecosystem, as they provide for the remediation of the habitats of its constituent species. These processes are vitally important, since water is regularly supplemented with autochthonous and allochthonous organic matter and biogenic elements brought by precipitation, fallout, and runoff from the adjoining territory. Thus, water self-purification for an aquatic ecosystem is an important function that ensures its stability, as is DNA repair for the system of heredity systems, which gives grounds for regarding water purification as ecological repair in aquatic ecosystems.

Table 2. Influence of different pollutants on the removal of suspended particles from water by filter-feeding hydrobionts. The effect on the efficiency of suspension removal (EESR) was calculated as described (Ostroumov, 2001a)

Pollutant	Hydrobiont	Concentration, mg/l	Note, reference
Trimethyl tin chloride (TMTC)	<i>Dreissena polymorpha</i>	0.01–10	Mitin, 1984; cited from Ostroumov, 2001a
Cadmium sulfate	<i>Mytilus galloprovincialis</i>	0.5	Original data
Copper sulfate	<i>M. galloprovincialis</i>	2	The same
Lead nitrate	<i>M. galloprovincialis</i>	20	»
Petroleum hydrocarbons (gas oil)	<i>M. galloprovincialis</i>	4–8	»
TDTMA	<i>M. edulis</i> × <i>M. galloprovincialis</i> (natural hybrid population)	0.05–5	»
TDTMA	<i>Crassostrea gigas</i>	0.5	EESR 761%
SDS	<i>M. edulis</i> , <i>M. galloprovincialis</i>	>1	Ostroumov, 2001a; Ostroumov, 2002a
SDS	<i>Crassostrea gigas</i>	0.5	EESR 231%
Triton X-100	<i>Unio tumidus</i>	5	Ostroumov, 2001a
Triton X-100	<i>M. edulis</i>	≥1	Ostroumov, 2001a
SD1 (OMO)	<i>Unio tumidus</i>	50	EESR 187%
SD2 (Tide)	<i>M. galloprovincialis</i>	50	EESR 207%
SD3 (Losk)	<i>M. galloprovincialis</i>	7	EESR 551%
SD4 (IXI)	<i>M. galloprovincialis</i>	10	EESR 158%
SD4 (IXI)	<i>M. galloprovincialis</i>	50	EESR 276%
LD1 (E)	<i>M. galloprovincialis</i>	2	EESR 214%
LD1 (E)	<i>Crassostrea gigas</i>	2	EESR 305%
LD2 (Fairy)	<i>Cr. gigas</i>	2	EESR 1790%

Note: The highest EESR value over the experimental period is indicated; SD is synthetic detergent, LD is liquid detergent, SDS is sodium dodecylsulfate.

RESPONSE OF THE SELF-PURIFICATION SYSTEM AS A WHOLE TO EXTERNAL (ANTHROPOGENIC) INFLUENCES ON A WATER BODY

The system of self-purification and water quality formation is labile (Ostroumov, 2000c) and easily transforms when environmental conditions change, which interferes with the analysis of trends in its functioning. The results of my experiments shed light on the factors responsible for the lability of a concrete process involved in self-purification, namely, water filtration by hydrobionts (mollusks and rotifers) (Ostroumov, 2000a–2000c, 2001a–2001c, 2002a–2002c, 2005). This process proved to be inhibited by sublethal concentrations of anthropogenic pollutants such as surfactants (Ostroumov *et al.*, 1997; Ostroumov, 2000a–2000c, 2001a), mixed surfactant-containing preparations (Ostroumov, 2001a; Ostroumov, 2005), cadmium, and some other substances (Table 2). A similar effect on mollusks and zooplanktonic filter-feeders was also described for other pollutants (Day and Kaushik, 1987). These data show the hazard of anthropogenic impact on aquatic ecosystems (chemical pollution of water bodies and water courses) as a factor impairing

the efficiency of the water self-purification system (Ostroumov, 2000a–c; 2001a–c; 2002a–c; Braginskii and Sirenko, 2003).

SOME GENERAL TENDENCIES AND PRINCIPLES OF FUNCTIONING OF THE WATER SELF-PURIFICATION SYSTEM

An analysis of data on specific features of the functioning of the biota as a factor of water self-purification in aquatic ecosystems provides a basis for some generalizations (note, however, that the tendencies listed below prevail but are not universal, and exceptions to the general rule may be found in some ecosystems):

(1) **The observed rates of particular self-purification processes are often lower than the maximum possible rates.** This may be evidence for the existence of regulatory mechanisms. Thus, the rate of water filtration by hydrobionts is regulated and considerably decreases when the concentration of suspension in water becomes higher (Sushchenya, 1975; Shul'man and Finenko, 1990).

(2) **There is a maximum diversification of organisms performing the main functions** in the mecha-

nism responsible for the parameters of the aquatic environment and its self-purification. Indeed, as noted above, almost every function (oxygen release, oxidation and transformation of dissolved organic matter, water filtration, etc.) in a given ecosystem is accomplished by different groups of organisms simultaneously.

(3) **The maximum possible number of stages in the pathways of biogenic migration of elements** is often characteristic of the functioning of biotic mechanisms responsible for the parameters of the aquatic environment and its purification.

(4) **Synecological cooperativity:** many processes that determine the parameters of the aquatic environment and lead to its self-purification are accomplished efficiently and at a high rate due to the joint actions of two or more species (groups of species).

(5) **Continuity of significance of the biota:** the biota remains highly important throughout the space occupied by an aquatic ecosystem and at any moment, irrespective of the time of day, season, or the stage of succession.

(6) **Balanced combination of opposite processes:** organisms simultaneously release and absorb organic molecules, oxygen, and CO₂; produce suspended organic matter and remove it from water in the course of their filtration (Ostroumov, 2002c); etc.

CONCLUSIONS AND RECOMMENDATIONS FOR NATURE-CONSERVATION PRACTICE

My experiments and the theoretical considerations based on their results suggest the following conclusions, which may be important for sustainable socioeconomic development:

(1) Since almost the entire aquatic biota is involved in the processes responsible for water quality and the self-purification of aquatic ecosystems or in their regulation, it is necessary to preserve the entire biodiversity of aquatic ecosystems (Ostroumov, 2002b).

(2) Since the species of terrestrial ecosystems and habitats bordering water bodies and watercourses actively participate in water purification, it is necessary to protect the biodiversity of these coastal ecosystems in order to maintain water quality at a high level.

(3) In estimating the critical anthropogenic load on an aquatic ecosystem (Moiseenko, 1999), it is important to take into account the lability and vulnerability of self-purification processes in the ecosystem.

(4) New types of ecological hazard created by chemical substances were revealed (Ostroumov, 2001c, 2002e).

(5) New principles were proposed for eutrophication control (Ostroumov, 2001b) and nature conservation in water areas (Ostroumov, 2002b).

(6) New pollutants that reduce the capacity of aquatic ecosystems (water bodies and watercourses) for self-purification will be revealed.

The multifunctional role of the biota in water self-purification is an additional illustration of V.I. Vernadsky's thesis that "... living matter ... geologically ... is the greatest force in the biosphere and determines all processes occurring in it" (Vernadsky, 1991).

ACKNOWLEDGMENTS

The author is grateful to A.P. Lisytsin, V.V. Malakhov, E.A. Kriksunov, G.S. Rozenberg, T.I. Moiseenko, V.D. Fedorov, V.V. Il'inskii, A.P. Sadchikov, O.F. Filenko, and G.E. Shul'man for discussing issues relevant to this study.

REFERENCES

- Alimov, A.F., Functional Ecology of Freshwater Mussels, *Tr. Zool. Inst. Akad. Nauk SSSR*, Leningrad, 1981, vol. 96.
- Alimov, A.F., *Elementy teorii funktsionirovaniya vodnykh ekosistem* (Principles of the Theory of Aquatic Ecosystem Functioning), St. Petersburg: Nauka, 2000.
- Braginskii, L.P. and Sirenko, L.A., Comprehensive Analysis of Toxic Hazard Created by Surfactants to Hydrobionts, *Gidrobiol. Zh.*, 2003, vol. 39, pp. 115–118.
- Dame, R., Bushek, D., and Prins, T., Benthic Suspension Feeders as Determinants of Ecosystem Structure and Function in Shallow Coastal Waters, *Ecol. Stud.*, 2001, vol. 151, pp. 11–37.
- Day, K. and Kaushik, N., Short-Term Exposure of Zooplankton to the Synthetic Pyrethroid, Fenvalerate, and Its Effects on Rates of Filtration and Assimilation of the Alga *Chlamydomonas reinhardtii*, *Arch. Environ. Contam. Toxicol.*, 1987, vol. 16, pp. 423–432.
- Filenko, O.F., *Vodnaya toksikologiya* (Aquatic Toxicology), Chernogolovka: Mosk. Gos. Univ., 1988.
- Izrael', Yu.A. and Tsyban', A.V., *Antropogennaya ekologiya okeana* (Anthropogenic Ecology of the Ocean), Leningrad: Gidrometeoizdat, 1989.
- Khailov, K.M., *Ekologicheskii metabolizm v more* (Ecological Metabolism in the Sea), Kiev: Naukova Dumka, 1971.
- Krasnoshchekov, G.P. and Rozenberg, G.S., Principles of Complication of Mechanisms Providing for Ecosystem Stability, in *Problemy ustoychivosti biologicheskikh sistem* (Problems of Biological System Stability), Moscow, 1992, pp. 40–51.
- Lisitsyn, A.P., Matter and Energy Fluxes in the External and Internal Earth Spheres, in *Global'nye izmeneniya prirodnoi sredy–2001* (Global Environmental Changes–2001), Dobretsov, N.L. and Kovalenko, V.I., Eds., Novosibirsk: Sib. Otd. Ross. Akad. Nauk, Fil. Geo, 2001, pp. 163–248.
- Matishov, D.G. and Matishov, G.G., *Radiatsionnaya ekologicheskaya okeanologiya* (Radiation–Ecological Oceanology), Apatity: Kol'sk. Nauch. Tsent. Ross. Akad. Nauk, 2001.
- Moiseenko, T.I., Methodology and Methods for Determining Critical Loads As Applied to Surface Waters of the Kola Subarctic Region, *Izv. Akad. Nauk, Ser. Geogr.*, 1999, no. 6, pp. 68–78.

- Monakov, A.V., *Pitanie presnovodnykh bespozvonochnykh* (Feeding of Freshwater Invertebrates), Moscow: IPEE, 1998.
- Ostroumov, S.A., *Vvedenie v biokhimicheskuyu ekologiyu* (Introduction to Biochemical Ecology), Moscow: Mosk. Gos. Univ., 1986.
- Ostroumov, S.A., Biological Filtering and Ecological Machinery for Self-Purification and Bioremediation in Aquatic Ecosystems: Towards a Holistic View, *Rivista di Biologia / Biology Forum*, 1998, vol. 91, pp. 221–232.
- Ostroumov, S.A., Aquatic Ecosystem: A Large Diversified Bioreactor with a Function of Water Self-Purification, *Dokl. Ross. Akad. Nauk*, 2000a, vol. 374, no. 3, pp. 427–429.
- Ostroumov, S.A., An Inhibitor Analysis of Regulatory Interactions in Trophic Chains, *Dokl. Ross. Akad. Nauk*, 2000b, vol. 375, no. 6, pp. 847–849.
- Ostroumov, S.A., The Concept of the Aquatic Biota as a Labile and Vulnerable Component in the System of Water Self-Purification, *Dokl. Ross. Akad. Nauk*, 2000c, vol. 372, no. 2, pp. 279–282.
- Ostroumov, S.A., *Biologicheskie efekty pri vozdeistvii pov-erkhnostno-aktivnykh veshchestv na organizmy* (Biological Effects of Surfactants), Moscow: MAKS-Press, 2001a.
- Ostroumov, S.A., Synecological Principles of Solving the Problem of Eutrophication, *Dokl. Ross. Akad. Nauk*, 2001b, vol. 381, no. 5, pp. 709–712.
- Ostroumov, S.A., Risk of Two-Level Synergism upon Synecological Summation of Anthropogenic Effects, *Dokl. Ross. Akad. Nauk*, 2001c, vol. 380, no. 6, pp. 847–849.
- Ostroumov, S.A., Inhibitory Analysis of Top-Down Control: New Keys to Studying Eutrophication, Algal Blooms, and Water Self-Purification, *Hydrobiologia*, 2002a, vol. 469, pp. 117–129.
- Ostroumov, S.A., Potentially Hazardous Substances with a New Type of Action: Disconnectors of the Pelagial–Benthic Interlink, *Dokl. Ross. Akad. Nauk*, 2002b, vol. 383, no. 1, pp. 138–141.
- Ostroumov, S.A., On the Role of Hydrobionts in Regulating the Fluxes of Matter and Migration of Elements in Aquatic Ecosystems, *Vestn. Ross. Akad. Est. Nauk*, 2002c, vol. 2, no. 3, pp. 50–54.
- Ostroumov, S.A., Conservation of Biodiversity and Water Quality: The Role of Feedback in Ecosystems, *Dokl. Ross. Akad. Nauk*, 2002d, vol. 382, no. 1, pp. 138–141.
- Ostroumov, S.A., Polyfunctional Role of Biodiversity in Processes Leading to Water Purification: Current Conceptualizations and Concluding Remarks, *Hydrobiologia*, 2002e, vol. 469, pp. 203–204.
- Ostroumov, S.A., A System of Principles for Preserving the Biogeocenotic Function and Biodiversity of Filter Feeders, *Dokl. Ross. Akad. Nauk*, 2002f, vol. 383, no. 5, pp. 710–713.
- Ostroumov, S.A., Concepts of Biochemical Ecology: Ecological Chemomediators, Ecological Chemoregulators, and Ecological Chemoeffectors, *Ecol. Stud. Probl. Solutions*, 2003, vol. 6, p. 105–107.
- Ostroumov, S.A., Some Aspects of Water Filtering Activity of Filter-Feeders, *Hydrobiologia*, 2005, vol. 542, pp. 275–286.
- Ostroumov, S.A., Donkin, P., and Stuff, P., An Anionic Surfactant Inhibits the Ability of *Mytilus edulis* Mussel to Filter and Purify Sea Water, *Vestn. Mosk. Gos. Univ., Ser. 16: Biol.*, 1997, no. 3, pp. 30–35.
- Ostroumov, S.A., Dodson, S., Hamilton, D., *et al.*, Medium-Term and Long-Term Priorities in Ecological Studies, *Rivista di Biologia / Biology Forum*, 2003, vol. 96, pp. 327–332.
- Sadchikov, A.P., Production and Transformation of Organic Matter by Size Groups of Phyto- and Bacterioplankton (An Example of Water Bodies of the Moscow Region), *Extended Abstract of Doctoral (Biol.) Dissertation*, Moscow, 1997.
- Shul'man, G.E. and Finenko, G.A., *Bioenergetika gidrobiontov* (Bioenergetics of Hydrobionts), Kiev: Naukova Dumka, 1990.
- Shvarts, S.S., Pyastolova, O.A., Dobrinskaya, L.A., and Runkova, G.G., *Effekt gruppy v populyatsiyakh vodnykh zivotnykh i khimicheskaya ekologiya* (The Group Effect in Aquatic Animal Populations and Chemical Ecology), Leningrad: Nauka, 1976.
- Skurlatov, Yu.I., Principles of Quality Control of Natural Waters, in *Ekologicheskaya khimiya vodnoi sredy* (Ecological Chemistry of the Aquatic Environment), Moscow, 1988, vol. 1, pp. 230–255.
- Sushchenya, L.M., *Kolichestvennye zakonomernosti pitaniya rakoobraznykh* (Quantitative Regularities of Crustacean Feeding), Minsk: Nauka i Tekhnika, 1975.
- Vernadsky, V.I., *Nauchnaya mysl' kak planetnoe yavlenie* (Scientific Thought as a Planetary Phenomenon), Moscow: Nauka, 1991.
- Vinogradov, M.E. and Shushkina, E.A., *Funktsionirovanie planktonnykh soobshchestv epipelagiali okeana* (The Functioning of Planktonic Communities in the Epipelagic Zone of the Ocean), Moscow: Nauka, 1987.
- Wetzel, R.G., *Limnology: Lake and River Ecosystems*, San Diego: Academic, 2001.
- Zavarzin, G.A. and Kolotilova, N.N., *Vvedenie v prirodovedcheskuyu mikrobiologiyu* (Introduction to Naturalist Microbiology), Moscow: Universitet, 2001.